

## Recombinant Human Interleukin-23/IL-23 (C-Fc)

**Catalog Number:** PKSH033874

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Interleukin-23; IL-23 protein Ile23-Ser328&Ala21-Pro189, with an C-terminal Fc
<b>Calculated MW</b>	81.3 kDa
<b>Observed MW</b>	80-110 kDa
<b>Accession</b>	P29460&Q9NPF7
<b>Bio-activity</b>	Measured by its ability to induce STAT reporter activity in 293F human embryonic kidney cells. The ED <sub>50</sub> for this effect is 179.42 ng/ml.


### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.


### Data



> 95 % as determined by reducing SDS-PAGE.



Measured by its ability to induce STAT reporter activity in 293F human embryonic kidney cells. The ED<sub>50</sub> for this effect is 179.42 ng/ml.



Immobilized Human IL-23R-His(PKSH032578) at 5µg/ml (100 µl/well) can bind Human IL-23-Fc (PKSH033874). The ED50 of Human IL-23-Fc (PKSH033874) is 0.187 µg/ml.

## Background

Interleukin 23 (IL-23) is a heterodimeric cytokine composed of two disulfide-linked subunits, a p19 subunit that is unique to IL-23, and a p40 subunit that is shared with IL-12. The p19 subunit has homology to the p35 subunit of IL-12, as well as to other single chain cytokines such as IL-6 and IL-11. The p40 subunit is homologous to the extracellular domains of the hematopoietic cytokine receptors. Although p19 is expressed by activated macrophages, dendritic cells, T cells, and endothelial cells, only activated macrophages and dendritic cells express p40 concurrently to produce IL-23. IL-23 has biological activities that are similar to, but distinct from IL-12. Both IL-12 and IL-23 induce proliferation and IFN-gamma production by human T cells. While IL-12 acts on both naive and memory human T cells, the effects of IL-23 is restricted to memory T cells.