

GNA12 Polyclonal Antibody

catalog number: **E-AB-90662**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human GNA12
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution

WB	1:500-1:2000
-----------	--------------

Data



Western blot analysis of extracts of various cell lines using GNA12 Polyclonal Antibody at 1:3000 dilution.

Observed-MW:44 kDa

Calculated-MW:34 kDa/37 kDa/44 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. Activates effector molecule RhoA by binding and activating RhoGEFs (ARHGEF12/LARG). GNA12-dependent Rho signaling subsequently regulates transcription factor AP-1 (activating protein-1) (By similarity). GNA12-dependent Rho signaling also regulates protein phosphatase 2A activation causing dephosphorylation of its target proteins. Promotes tumor cell invasion and metastasis by activating RhoA/ROCK signaling pathway and up-regulating proinflammatory cytokine production. Inhibits CDH1-mediated cell adhesion in process independent from Rho activation. Together with NAPA promotes CDH5 localization to plasma membrane. May play a role in the control of cell migration through the TOR signaling cascade.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8