# **Elabscience**®

### Human MSP Antibody Pair Set

Catalog No.E-KAB-0434ApplicationsELISASynonymsMST1;D3F15S2;DNF15S2;HGFL;MSP;NF15S2;macrophage stimulating 1;Hepatocyte growth<br/>factor-like protein;Macrophage stimulatory protein;Macrophage-stimulating protein

#### **Kit components & Storage**

Title	Specifications	Storage
Human MSP Capture Antibody	1 vial, 100 µ g	Store at $-20^{\circ}$ C for one year. Avoid
		freeze/thaw cycles.
Human MSP Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year. Avoid
		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0434)	
		Human MSP Capture Antibody	Human MSP Detection Antibody (Biotin)
Immunogen	Immunogen	Recombinant Human MSP protien	Recombinant Human MSP protien
Information	Swissprot	P26927	
Product details	Reactivity	Human	Human
	Host	Mouse	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Protein A or G	Protein A or G
	Specificity	Detects Human MSP in ELISAs.	

For Research Use Only

## **Elabscience**®

#### Applications

Human MSP Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human MSP Capture	
Capture		Antibody	10
			A result of the second se
ELISA	1:1000-1:10000	Human MSP Detection	Optical Density
Detection		Antibody (Biotin)	•
			0. 01
			Human MSP concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

#### Background

The protein encoded by this gene contains four kringle domains and a serine protease domain, similar to that found in hepatic growth factor. Despite the presence of the serine protease domain, the encoded protein may not have any proteolytic activity. The receptor for this protein is RON tyrosine kinase, which upon activation stimulates ciliary motility of ciliated epithelial lung cells. This protein is secreted and cleaved to form an alpha chain and a beta chain bridged by disulfide bonds.

For Research Use Only