A Reliable Research Partner in Life Science and Medicine

Recombinant Human Interleukin-2/IL-2 Protein (His Tag)

Catalog Number: PKSH033425

Note: Centrifuge before opening to ensure complete recovery of vial contents.

T.			
	COL	TM	tion
D.C.	201	ци	ш

Species Human

Source E.coli-derived Human Interleukin-2/IL-2 protein Ala21-Thr153, with an C-terminal His

Calculated MW16.4 kDaObserved MW13 kDaAccessionP60568

Bio-activity Measure by its ability to induce proliferation in CTLL-2 cells. The ED_{50} for this effect

is <3 ng/mL. The specific activity of recombinant human IL-2 is >5 x 10^6 IU/mg.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.01 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS,pH 8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.