

Vimentin Polyclonal Antibody

catalog number: AN005800L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human Vimentin protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% Proclin300, 1% protective protein and 50% glycerol, pH7.4

Applications Recommended Dilution

WB	1:500-1:1000
-----------	--------------

Data



Western blot with Anti Vimentin Polyclonal antibody at dilution of 1:1000. Lane 1: U-2 OS cell lysate, Lane 2: 293 cell lysate.

Observed-MW:57 kDa

Calculated-MW:54 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Vimentin is a type III intermediate filament (IF) protein found in various non-epithelial cells, especially mesenchymal cells. A vimentin monomer, has a central α -helical domain and carboxyl (tail) domains. Two monomers compose the basic subunit of vimentin assembly. Vimentin is crucial for supporting and anchoring the position of the organelles in the cytosol. Vimentin provided cells with a resilience absent from the microtubule or actin filament networks, when under mechanical stress in vivo. Therefore, in general, it is accepted that vimentin is the cytoskeletal component responsible for maintaining cell integrity. Vimentin is also responsible for stabilizing cytoskeletal interactions. It is found that vimentin control the transport of low-density lipoprotein. It has been used as a sarcoma tumor marker to identify mesenchyme.

For Research Use Only