# APC Anti-Human CD45 Antibody[QA17A19]

**Catalog Number:** AN00358E

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Clone No.** | QA17A19 |
| **Isotype Control** | APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E] |
| **Conjugation** | APC |
| **Conjugation Information** | APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter). |
| **Storage Buffer** | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. |

| Applications | Recommended usage |
|---|---|
| **FCM** | Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. |

## Preparation & Storage

| | |
|---|---|
| **Storage** | Keep as concentrated solution.<br>This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze. |
| **Shipping** | Ice bag |

## Antigen Information

| | |
|---|---|
| **Alternate Names** | Leukocyte Common Antigen;LCA;T200 |
| **Uniprot ID** | P08575 |
| **Gene ID** | 5788 |
| **Background** | CD45 is a 180 - 240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes or platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPA P) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4 . |


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com