APC Anti-Human IL-12/IL-23 p40 Antibody[C8.6]

Catalog Number: AN00843E

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	C8.6
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications — Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



IFN-γ-primed and then LPS-stimulated (24h) human peripheral blood monocytes intracellular stained with APC Anti-Human IL-12/IL-23 p40 Antibody[C8.6] (filled gray histogram) or APC Mouse IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Interleukin-1α, Lymphocyte-activating factor (LAF), Endogenous pyrogen (EP), Leukocyte endogenous mediator (LEM), Mononuclear cell factor (MCF/MNCF)
Uniprot ID	P29460

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com

Gene ID	3593
Background	The C8.6 antibody reacts with human IL-12 p40 subunit of the IL-12 p70 heterodimer and IL-23 p40 subunit of the IL-23 p19/p40, as well as p40 monomer and homodimer. The C8.6 antibody has been reported to strongly inhibit different biological activities of IL-12, (e.g., IFN-gamma induction, mitogenic effects on PHA blasts, and enhancement of NK cell-mediated cytotoxicity). The C8.6 antibody can neutralize the bioactivity of natural or recombinant IL-12.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com