


## TRIP4 Polyclonal Antibody

catalog number: E-AB-14750

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Mouse
Immunogen	Recombinant protein of human TRIP4
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:25-1:100

## Data




Western Blot analysis of A431 cell using TRIP4 Polyclonal Antibody at dilution of 1:400

Calculated-MV:66 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer using TRIP4 Polyclonal Antibody at dilution of 1:50



Immunohistochemistry of paraffin-embedded Human liver cancer using TRIP4 Polyclonal Antibody at dilution of 1:50

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

Activating signal co-integrator-1 (ASC-1), originally identified as TRIP4, is a transcriptional co-activator of nuclear receptors that associates with specific components of the RNA polymerase II complex and binds the basal transcription factors TBP and TFIIB. ASC-1 functions with the transcription integrators SRC-1 and CBP/p300 through its zinc finger motif and is dependent on their ligand-dependent transactivation domain, AF2. Endogenous ASC-1 in HeLa cells is predominantly a nuclear protein. Under conditions of serum starvation, ASC-1 localizes to the cytoplasm.

## For Research Use Only