



A Reliable Research Partner in Life Science and Medicine

Recombinant S100A12 Monoclonal Antibody

catalog number: AN300131P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human S100A12 protein


HostRabbitIsotypeIgGCloneA1245PurificationProtein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

IHC-P 1:100-1:500

Data



Immunohistochemistry of paraffin-embedded human tonsil using S100A12 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20 $^{\circ}\text{C}$ to -80 $^{\circ}\text{C}$. Preservative-Free. Avoid repeated freeze-thaw cycles .

Shipping Ice bag

Background

S100 protein is a family of low molecular weight protein found in vertebrates characterized by two EF-hand calciumbinding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is 100% soluble in ammonium sulfate at neutral pH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from the neural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A12, also known as S100 calcium-binding protein A12, Calcium-binding protein in amniotic fluid 1, Calgranulin-C, and S100A12, is a member of the S-101 family. Like the majority of S100 proteins, S100A12 is a dimer, with the interface between the two subunits being composed mostly of hydrophobic residues. The fold of S100A12 is similar to the other known crystal and solution structures of S100 proteins, except for the linker region between the two EF-hand motifs. S100A12 plays an important role in the inflammatory response.

For Research Use Only