

## Purified Anti-Mouse/Rat CD29 Antibody[HMβ1-1]

catalog number: E-AB-F1309A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse/Rat
<b>Immunogen</b>	Recombinant Mouse CD29 protein
<b>Host</b>	Armenian hamster
<b>Isotype</b>	Armenian hamster IgG
<b>Clone</b>	HMβ1-1
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications

### Recommended Dilution

<b>FCM</b>	2 µg/mL(0.5×10 <sup>6</sup> -1×10 <sup>6</sup> cells)
------------	---

### Data



Rat bone marrow lymphocytes were stained with 0.2 µg Purified Anti-Mouse/Rat CD29 Antibody[HMβ1-1] (Right) and 0.2 µg Armenian Hamster IgG, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody, then anti-Rat CD19 PE-conjugated Monoclonal Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

CD29 is a 130 kDa protein, also known as integrin β1, VLA-β chain, or GPIIa. It is a member of the integrin family, expressed broadly on leukocytes, endothelial cells, smooth muscle, and epithelial cells. In association with CD49a-f, CD29 forms the VLA-1 through VLA-6 complexes, respectively. It plays an important role in cell-cell or cell-matrix interaction.

### For Research Use Only