

Recombinant Human NCAM1 Protein (His Tag)

Catalog Number: PKSH032793

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human NCAM1 protein Leu20-Pro603, with an C-terminal His
Calculated MW	65.5 kDa
Observed MW	90-120 kDa
Accession	P13591-3
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Neural cell adhesion molecule 1 (NCAM-1) is a single-pass type I membrane protein, it belongs to a family of membrane-bound glycoproteins that are involved in Ca²⁺ independent cell matrix and homophilic or heterophilic cell-cell interactions. NCAM-1 is synthesized as a 761 aa preproprecursor that contains a 19 aa signal sequence, a 722 aa GPI-linked mature region, and a 20 aa C-terminal prosegment. The molecule contains five C-2 type Ig-like domains and two fibronectin type-III domains. NCAM-1 is a cell adhesion molecule involved in neuron-neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. Acting as a receptor for rabies virus, NCAM-1 in the adult brain shows a decline of sialylation relative to earlier developmental periods.

For Research Use Only