Elabscience®

SLC16A1 Polyclonal Antibody


catalog number: E-AB-15768


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Synthetic peptide of human SLC16A1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Pacammandad Dilution

ApplicationsRecommended DilutionIHC1:50-1:200

Data





Immunohistochemistry of paraffin-embedded Human liver Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using SLC16A1 Polyclonal Antibody at dilution cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilution 1:50 cancer tissue using SLC16A1 Polyclonal Antibody at dilut

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

The protein encoded by this gene is a proton-linked monocarboxylate transporter that catalyzes the movement of many monocarboxylates, such as lactate and pyruvate, across the plasma membrane. Mutations in this gene are associated with erythrocyte lactate transporter defect. Alternatively spliced transcript variants have been found for this gene.