

A Reliable Research Partner in Life Science and Medicine

Recombinant Human Cathepsin E/CTSE Protein (His Tag)

Catalog Number: PKSH032181

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip					
	00	CI	411	n	П	n	m
JU	\mathbf{c}	U		J A	ш	v.	ш

Species Human

Source HEK293 Cells-derived Human Catheps in E; CTSE protein Ser20-Pro396, with an C-

terminal His

Calculated MW41.8 kDaObserved MW46 kDaAccessionP14091

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution of 20mM MES, 150mM NaCl, pH 5.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cathepsin E (CTSE) is a gastric aspartyl protease that functions as a disulfide-linked homodimer. It is a member of the Peptidase C1 family, and has a specificity similar to that of Pepsin A and Cathepsin D. CTSE is localized to the endoplasmic reticulum and Golgi apparatus, while the mature enzyme is localized to the endosome. It is expressed abundantly in the stomach, the Clara cells of the lung and activated B-lymphocytes, and at lower levels in lymph nodes, skin and spleen. CTSE is an intracellular proteinase that have a role in immune function, activation-induced lymphocyte depletion in the thymus, neuronal degeneration and glial cell activation in the brain. Futhermore, it probably involved in the processing of antigenic peptides during MHC class II-mediated antigen presentation.

For Research Use Only