

Recombinant Human Interleukin-17F/IL-17F Protein

Catalog Number: PKSH032622

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human Interleukin-17F;IL-17F protein Arg31-Gln163, with an C-terminal His
Calculated MW	15.8 kDa
Observed MW	18 kDa
Accession	Q96PD4
Bio-activity	Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED ₅₀ for this effect is <20 ng/mL.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM sodium acetate, pH 4.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Interleukin-17 is a potent pro-inflammatory cytokine produced by activated memory T cells. There are at least six members of the IL-17 family in humans and in mice. Today, IL-17 represents a family of structurally related cytokines that share a highly conserved C-terminal region but differ from one another in their N-terminal regions and in their distinct biological roles. The six known members of this family; IL-17A through IL-17F; are secreted as homodimers. IL-17F also regulates cartilage matrix turnover and inhibits angiogenesis.

For Research Use Only