

A Reliable Research Partner in Life Science and Medicine

# **HSP40-4 Polyclonal Antibody**

catalog number: E-AB-11313

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant protein of human DNAJA1

Host Rabbit
Isotype IgG

**Purification** Affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

## **Applications** Recommended Dilution

**WB** 1:500-1:2000 **IHC** 1:100-1:300


#### Data




Immunohistochemistry of paraffin-embedded Human colon

Western Blot analysis of HepG2, Raji, A431 and 231 cell using HSP40-4 Polyclonal Antibody at dilution of 1:800

cancer using HSP40-4 Polyclonal Antibody at dilution of





Immunohistochemistry of paraffin-embedded Human thyroid cancer using HSP40-4 Polyclonal Antibody at dilution of 1:60

# Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

# Background

#### For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Fax: 1-832-243-6017

## **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

**Elabscience®** 

DnaJ-like proteins interact with HSP 70 molecular chaperones and function to facilitate protein folding and mitochondrial protein import. HSP 40-4, also known as HDJ2, is the human DnaJ homolog that functions as a co-chaperone with a cysteine-rich zinc finger domain. The cellular redox enzyme thioredoxin interacts with HSP 40-4, and oxidation and reduction reversibly regulate HSP 40-4 function in response to the changing redox states of the cell. The zinc finger domain of HSP 40-4 may act as a redox sensor of chaperone-mediated protein-folding machinery, since HSP 40-4 inactivation leads to the oxidation of cysteine thiols and a simultaneous release of coordinated zinc. Loss of the HSP 40-4 protein may be linked to severe defects in spermatogenesis that involve aberrant androgen signaling.

Fax: 1-832-243-6017