

PE/Elab Fluor® 594 Anti-Mouse CD25 Antibody[PC-61.5.3]

Catalog Number: E-AB-F1102UP

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

| | |
|-------------------------|---|
| Reactivity | Mouse |
| Host | Rat |
| Isotype | Rat IgG1, κ |
| Clone No. | PC-61.5.3 |
| Isotype Control | PE/Elab Fluor® 594 Rat IgG1, κ Isotype Control[HRPN] [Product E-AB-F09823P] |
| Conjugation | PE/Elab Fluor® 594 |
| Conjugation Information | PE/Elab Fluor® 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA. |


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶ cells in 100 µL volume].

Data



C57BL/6 murine splenocytes are stained with APC Anti-

Mouse CD4 Antibody and PE/Elab Fluor® 594 Anti-Mouse CD25 Antibody[PC-61.5.3] (Left). Splenocytes are stained

with APC Anti-Mouse CD4 Antibody and PE/Elab Fluor® 594 Rat IgG1, κ Isotype Control (Right).

Preparation & Storage

Storage

Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.

Shipping

Ice bag

Antigen Information

Alternate Names

IL-2 receptor subunit alpha;IL-2-RA;IL-2R subunit alpha;IL2-RA;IL2RA;Interleukin-2 receptor subunit alpha;TAC antigen;p55

For Research Use Only

Uniprot ID

P01590

Gene ID

16184

Background

CD25 is a 55 kD glycoprotein, also known as the low affinity IL-2R α , Ly-43, p55, or Tac. It is expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2R β) and CD132(common γ chain), CD25 forms the high affinity signaling IL-2 receptor.