Elabscience®

CDK4 Polyclonal Antibody

catalog number: E-AB-13989

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant protein of human CDK4
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
Data	
	$ \frac{k_{0}}{k_{0}} = \frac{1}{2} $ $ \frac{1}{2} = \frac{1}{2} $ $ \frac{1}{2} = \frac{1}{2} $ $ \frac{1}{2} = \frac{1}{2} $
•	is of Hela and 293T cell using CDK4
Polyclonal	Antibody at dilution of 1:450
Cal	culated-MW:34 kDa
Preparation & Stora	ge
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

C 1	
Shipping	

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the protein kinase complex that is important for cell cycle Gl phase progression. The activity of this kinase is restricted to the Gl-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and Rb were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation sites of this gene have been reported.