
BMI1 Polyclonal Antibody


Catalog Number:E-AB-64659



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|---|
| Reactivity | Human |
| Immunogen | Recombinant fusion protein of human BMI1 (NP_005171.4). |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Affinity purification |
| Conjugation | Unconjugated |
| Formulation | PBS with 0.02% sodium azide, 50% glycerol, pH7.3. |
| Applications | Recommended Dilution |
| WB | 1:500-1:2000 |
| IF | 1:50-1:200 |
| Data | |





Western blot analysis of extracts of U-937 cells using BMI1 Polyclonal Antibody at dilution of 1:1000. Observed Mw:41kDa Calculated Mw:36kDa

Immunofluorescence analysis of HeLa cells using BMI1 Polyclonal Antibody

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a ring finger protein that is major component of the polycomb group complex 1 (PRC1). This complex functions through chromatin remodeling as an essential epigenetic repressor of multiple regulatory genes involved in embryonic development and self-renewal in somatic stem cells. This protein also plays a central role in DNA damage repair. This gene is an oncogene and aberrant expression is associated with numerous cancers and is associated with resistance to certain chemotherapies. A pseudogene of this gene is found on chromosome X. Read-through transcription also exists between this gene and the upstream COMM domain containing 3 (COMMD3) gene.

For Research Use Only

A Reliable Research Partner in Life Science and MedicineToll-free: 1-888-852-8623Tel: 1-832-243-6086Web: www.elabscience.comEmail: techsupport@elabscience.com