

## Recombinant PEDF Monoclonal Antibody

catalog number: **AN301984L**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

### Description


|                     |                                                                 |
|---------------------|-----------------------------------------------------------------|
| <b>Reactivity</b>   | Human                                                           |
| <b>Immunogen</b>    | Peptide. This information is proprietary to PTMab.              |
| <b>Host</b>         | Rabbit                                                          |
| <b>Isotype</b>      | Rabbit IgG                                                      |
| <b>Clone</b>        | C331605                                                         |
| <b>Purification</b> | Protein A purified                                              |
| <b>Buffer</b>       | PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant. |

### Applications

### Recommended Dilution

|           |               |
|-----------|---------------|
| <b>WB</b> | <b>1:1000</b> |
|-----------|---------------|

### Data



Western Blot with PEDF Monoclonal Antibody at dilution of 1:1000. Lane 1: K562, Lane 2: TF-1

**Observed-MW:46 kDa**

**Calculated-MW:46 kDa**

### Preparation & Storage

|                 |                                                                 |
|-----------------|-----------------------------------------------------------------|
| <b>Storage</b>  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| <b>Shipping</b> | Ice bag                                                         |

### Background

PEDF also known as serpin F1 (SERPINF1), is a multifunctional secreted protein that has anti-angiogenic, anti-tumorigenic, and neurotrophic functions. It is an approximately 50-kDa secreted protein that is widely expressed, including by osteoblasts and osteoclasts. PEDF is also one of the most abundant secretory products of adipocytes, and circulating concentrations of PEDF correlate positively with body fat mass and INS resistance. PEDF is a neurotrophic factor involved in neuronal differentiation in retinoblastoma cells. Mutations in this gene were found in individuals with osteogenesis imperfecta, type VI.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.2