

A Reliable Research Partner in Life Science and Medicine

FITC-conjugated Myc-Tag Monoclonal Antibody

catalog number: AN00300C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}			4.0	
- 1	ec.	cri	пπ	OH
ய	CO.	CIL	JUL	UL

Reactivity All

Immunogen MYC tag fusion protein


Host Mouse Isotype IgGl Clone 6B10

PurificationProtein A/G PurificationConjugationFITC(Ex/Em:495nm/525nm)

Buffer PBS with 0.05% Proclin300 and 0.1% gelatin, pH7.4

FCM $0.2 \mu g \text{ per } 10^6 \text{ cells in a } 100 \mu L \text{ suspension}$

Data



1X10^6 HEK-293F cells were intracellularly stained with 0.2 ug FITC-conjugated Myc-Tag Monoclonal Antibody (red), or 0.2 ug FITC-conjugated Myc-Tag Monoclonal Antibody Isotype Control (green).

Preparation & Storage

Storage Storage Store at 4°C Valid for 12 months. Avoid freeze / thaw cycles. Protected from prolonged

exposure to light.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombin

Web: www.elabscience.cn