# Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

# Recombinant Human IL-3 Receptor Subunit Alpha/IL-3RA/CD123 (C-6His)

Catalog Number: PKSH034052

Note: Centrifuge before opening to ensure complete recovery of vial contents.

ъ.					7.6		
IDZ	e s	C	r٦	n	Ħ	n	П

Species Human

Source HEK293 Cells-derived Human IL-3RA; CD123 protein Thr19-Arg305, with an C-terminal

His

 Mol\_Mass
 33.9 kDa

 Accession
 P26951

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

CD123, also known as Interleukin-3 receptor subunit alpha, belongs to the type I cytokine receptor family. In mouse, there are two classes of high-affinity IL3 receptors. One contains an IL3-specific beta subunit and the other contains the beta subunit also shared by high-affinity IL5 and GM-CSF receptors. CD123 stimulates the proliferation and differentiation of hemopoietic cells including the pluripotent hematopoietic stem cells as well as various lineage-committed cells. CD123 is a heterodimer consisting of an alpha and a beta subunit. The alpha subunit alone binds IL-3 with low affinity. The beta subunit does not bind IL-3 by itself but is required for the high-affinity binding of IL-3 to the heterodimeric receptor complex.

#### For Research Use Only