

Recombinant Human IL-1R9/IL1RAPL2 Protein (Fc Tag)

Catalog Number: PKSH031827

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Human |
| Source | HEK293 Cells-derived Human IL-1R9/IL1RAPL2 protein Met 1-Glu 356, with an C-terminal hFc |
| Calculated MW | 66.0 kDa |
| Observed MW | 80-85 kDa |
| Accession | NP_059112.1 |
| Bio-activity | Measured by its ability to bind biotinylated human IL1 α in functional ELISA. |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

X-linked interleukin-1 receptor accessory protein-like 2 (IL1RAPL2) or Interleukin-1 receptor 9 (IL-1R9) is a member of the interleukin 1 receptor family. This protein is similar to the interleukin 1 accessory proteins. IL-1R9/IL1RAPL2 shows restricted expression in fetal brain and is highly homologous to IL1RAPL; which is reportedly involved in nonsyndromic X-linked mental retardation. IL-1R9/IL1RAPL2 is highly homologous to IL-1R8. Both forms have no known ligands and receptor are found in the fetal brain. IL-1R9/IL1RAPL2 may function as a negative receptor. Both IL1RAPL1 and IL1RAPL2 have novel C-terminal sequences not present in other related proteins. IL-1R9/IL1RAPL2 may be strong candidates for X-linked non-syndromic mental retardation loci; and that molecules resembling IL-1 and IL-18 play a role in the development or function of the central nervous system.