

## Recombinant Human TNFRSF19/TROY Protein (His Tag)

**Catalog Number:** PKSH030638

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human TNFRSF19/TROY protein Met 1-Leu170, with an C-terminal His
<b>Calculated MW</b>	16.9 kDa
<b>Observed MW</b>	28 kDa
<b>Accession</b>	Q9NS68-2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Tumor necrosis factor receptor superfamily; member 19 (TNFRSF19); also known as TAJ-alpha or TROY; is a member of the TNF-receptor superfamily. TNFRSF19/TROY expression is detected in the pulmonary epithelium and the ductal epithelium of the prostate and parotid glands. TNFRSF19/TROY expression is detected in some adenocarcinoma cell lines that arise from this tissue. It has been shown to interact with TRAF family members; and to activate JNK signaling pathway when overexpressed in cells. TNFRSF19/TROY is capable of inducing apoptosis by a caspase-independent mechanism; and it is thought to play an essential role in embryonic development. TNFRSF19/TROY was negatively regulated by adipogenic transcription factor CCAAT/enhancer-binding proteins (C/EBP). TNFRSF19 signals activation of the Jnk pathway and induces cell death. Overexpression of TNFRSF19 also signals NFB activation; comparable and similar to that by p75NGFR. TNFRSF19/TROY is capable of activating key signaling pathways of the TNF receptor family; and its predominant expression patterns suggest that it plays a role in the growth and regulation of epithelial tissues.

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017