

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 700 Anti-Mouse CD11c Antibody[N418]

Catalog Number: E-AB-F0991M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Host Armenian Hamster
Isotype Armenian Hamster IgG

Clone No. N418

Isotype Control Elab Fluor® 700 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852M1]

Conjugation Elab Fluor® 700

Conjugation Information Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse MHC II (I-A/I-E) Antibody[M5/114] and Elab Fluor[®] 700 Anti-Mouse CD11c Antibody[N418] (left) or Elab Fluor[®] 700 Armenian Hamster IgG Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion

receptor p150+95

Uniprot ID Q9QXH4

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 16411

CD11c is a 150 kD glycoprotein also known as α X integrin, CR4, and p150. CD11c forms a α X β 2 heterodimer with β 2 integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The α X β 2 integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.