Elabscience®


Elab Fluor[®] 700 Anti-Mouse CD11c Antibody[N418]

Catalog Number: E-AB-F0991M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone No.	N418
Isotype Control	Elab Fluor [®] 700 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852M1]
Conjugation	Elab Fluor [®] 700
Conjugation Information	Elab Fluor [®] 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse MHC II (I-A/I-E) Antibody[M5/114] and Elab Fluor[®] 700 Anti-Mouse CD11c Antibody[N418] (left) or Elab Fluor[®] 700 Armenian Hamster IgG Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage	•
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD11 antigen-like family member C;CD11c;Integrin alpha-X;Itgax;Leukocyte adhesion receptor p150+95

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>www.elabscience.com</u>

Elabscience®

Uniprot ID Gene ID Background Q9QXH4

16411

CD11c is a 150 kD glycoprotein also known as α X integrin, CR4, and p150. CD11c forms a α X β 2 heterodimer with β 2 integrin (CD18). It is primarily expressed on dendritic cells, NK cells, a subset of intestinal intraepithelial lymphocytes (IEL), and some activated T cells. The α X β 2 integrin plays an important role in cell-cell contact by binding its ligands: iC3b, fibrinogen and CD54.