

Recombinant Cynomolgus IL-3 Receptor Subunit Alpha/IL-3RA/CD123 (C-Fc)

Catalog Number: PKSQ050122

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Cynomolgus macaques |
| Source | HEK293 Cells-derived Cynomolgus macaques IL-3RA/CD123 protein Arg18-Arg302, with an C-terminal His |
| Calculated MW | 33.6 kDa |
| Observed MW | 50-65 kDa |
| Accession | G8F3K3 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

CD123, also known as Interleukin-3 receptor subunit alpha, belongs to the type I cytokine receptor family. In mouse, there are two classes of high-affinity IL3 receptors. One contains an IL3-specific beta subunit and the other contains the beta subunit also shared by high-affinity IL5 and GM-CSF receptors. CD123 stimulates the proliferation and differentiation of hemopoietic cells including the pluripotent hematopoietic stem cells as well as various lineage-committed cells. CD123 is a heterodimer consisting of an alpha and a beta subunit. The alpha subunit alone binds IL-3 with low affinity. The beta subunit does not bind IL-3 by itself but is required for the high-affinity binding of IL-3 to the heterodimeric receptor complex.