

Recombinant Mouse FKBP12 Protein (His Tag)

Catalog Number: PKSM040771

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	E.coli-derived Mouse FKBP12 protein Gly 2-Glu 108, with an N-terminal His
Calculated MW	12.7 kDa
Accession	NP_032045.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

FK506 binding protein 12 (FKBP12), also known as FKBP1, along with cyclophilin, are two major members of the immunophilin protein family who serve as receptors for the immunosuppressant drugs cyclosporin A and FK506. FKBP12 has attracted immense attention and its role in mediating the immunosuppressive functions. FKBP12 serves a dual role as a peptidyl-prolyl cis-trans isomerase and as a modulator of several cell signaling pathways. In one such a role, FKBP12 interacts with and regulates the functional state of the ryanodine Ca²⁺ channel receptor by altering protein conformation and coordinating multi-protein complex formation. Another physiological role of FKBP12 is an interactor and a regulator of the type I serine/threonine kinase receptors of TGF-beta superfamily. Current data, derived from detailed biochemical studies as well as from functional studies in various systems, suggest that FKBP12 functions as a guardian, for the type I receptors to prevent them from leaky signaling under sub-optimal ligand concentrations, thereby providing a molecular gradient reader, for TGF-beta family morphogens. This aspect of FKBP12 function may be critical for cellular responsiveness to morphogenetic gradients of the TGF-beta family members during early development, serving to assure the translation of different ligand concentrations into different signaling readouts. In addition, FKBP12 may be involved in neuronal or astrocytic cytoskeletal organization and in the abnormal metabolism of tau protein in Alzheimer's disease (AD) damaged neurons.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017