Recombinant Human NETO1/BTCL1 Protein (His Tag)

Catalog Number: PKSH031084

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human NETO1/BTCL1 protein Met 1-Thr 344, with an C-
	terminal His
Calculated MW	38 kDa
Observed MW	46 kDa
Accession	NP_620416.1
Bio-activity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Elabscience®

Neuropilin tolloid-like 1 (NETO1), a complement Clr/Cls, Uegf, Bmp1 (CUB) domain-containing transmembrane protein, is a novel component of the NMDAR complex critical for maintaining the abundance of NR2A-containing NMDARs in the postsynaptic density. The N-methyl-D-aspartate receptor (NMDAR), a major excitatory ligand-gated ion channel in the central nervous system (CNS), is a principal mediator of synaptic plasticity. Both NETO1 and NETO2 share an identical and unique domain structure thus representing a novel subfamily of CUB- and LDLa-containing proteins. The cytoplasmic domains of NETO1 and NETO2 are not homologous to other known protein sequences but contain a conserved FXNPXY-like motif, which is essential for the internalization of clathrin coated pits during endocytosis or alternatively, may be implicated in intracellular signaling pathways. NETO1 and NETO2, have marked effects on receptor properties, increasing further the potential diversity of Kainate receptors (KARs) functional properties. NETO1 involves in the development and/or maintenance of neuronal circuitry. NETO1 regulates long-term NMDA receptor-dependent synaptic plasticity and cognition, at least in the context of spatial learning and memory.