Elabscience®


Elab Fluor[®] Violet 450 Anti-Human CD14 Antibody[UCHM-1]

Catalog Number: AN00420Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse IgG2a, к
Clone No.	UCHM-1
Isotype Control	Elab Fluor [®] Violet 450 Mouse IgG2a, к Isotype Control[C1.18.4] [Product E-AB- F09802Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.





Staining of normal human peripheral blood cells with Elab

Fluor[®] Violet 450 Anti-Human CD14 Antibody[UCHM-1] (filled gray histogram) or Elab Fluor[®] Violet 450 Mouse IgG2a, κ Isotype Control (empty black histogram). Cells in the monocytes gate were used for analysis.

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	Monocyte differentiation antigen CD14;myeloid cell-specific leucine-rich glycoprotein;
	LPS receptor
For Research Use Only	

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Elabscience®

Uniprot ID	
Gene ID	
Background	

P08571

929

CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein that is also known as the LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens and in the upregulation of adhesion molecules and cytokine expression in monocytes and neutrophils.