

## Recombinant Rat DDR2 Kinase/CD167b Protein (Fc Tag)

**Catalog Number:** PKSR030218

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	HEK293 Cells-derived Rat DDR2 Kinase/CD167b protein Met1-Arg399, with an C-terminal hFc
<b>Calculated MW</b>	69.5 kDa
<b>Observed MW</b>	110 kDa
<b>Accession</b>	B1WC09
<b>Bio-activity</b>	Measured by its ability to bind with Rat tail Collagen I in a functional ELISA.

### Properties

<b>Purity</b>	> 80 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 80 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Discoidin domain receptor 2 (DDR2) or CD167b (cluster of differentiation 167b) is a kind of protein tyrosine kinases associated with cell proliferation and tumor metastasis, and collagen, identified as a ligand for DDR2, up-regulates matrix metalloproteinase 1 (MMP-1) and MMP-2 expression in cellular matrix. DDR2/CD167b was found to recognise the triple-helical region of collagen X as well as the NC1 domain. Binding to the collagenous region was dependent on the triple-helical conformation. DDR2/CD167b autophosphorylation was induced by the collagen X triple-helical region but not the NC1 domain, indicating that the triple-helical region of collagen X contains a specific DDR2 binding site that is capable of receptor activation. DDR2/CD167b is induced during stellate cell activation and implicate the phosphorylated receptor as a mediator of MMP-2 release and growth stimulation in response to type I collagen. Moreover, type I collagen-dependent upregulation of DDR2/CD167b expression establishes a positive feedback loop in activated stellate cells, leading to further proliferation and enhanced invasive activity.

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017