

Purified Anti-Human TCRV γ 9 Antibody[B3]

catalog number: AN003570P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human TCRV γ 9 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	B3
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
buffer	PBS, pH 7.2. Contains 0.05% proclin 300.

Applications

Recommended Dilution

FCM	2 μ g/mL (1×10^5 - 5×10^5 cells)
------------	---

Data



Human peripheral blood lymphocytes were stained with 0.2 μ g Purified Anti-Human TCRV γ 9 Antibody[B3] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The V γ 9 TCR is a variant of the TCR γ chain expressed on a subset of γ/δ T cells. V γ 9V δ 2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomeric proteins CD3 δ , γ , ϵ , and ζ chains. The B3 antibody reacts specifically with human TCR V γ 9 as designated by the Lefranc/Foster nomenclature system. Human TCR V γ 9 is also known as TCR V γ 2 under the Strauss, Quertermous nomenclature system. Several TCR γ and δ chain nomenclature systems exist. In order to consolidate the various nomenclature systems, we have created charts for corresponding names across some of the most popular naming methods

For Research Use Only