Elabscience®

CD218b Polyclonal Antibody

catalog number: E-AB-14440

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Applications Recommended Dilut WB 1:500-1:2000 Data Image: I	n, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Host Rabbit Isotype IgG Purification Affinity purification Buffer Phosphate buffered solution Applications Recommended Dilut WB 1:500-1:2000 Data Data Western Blot analysis of RAW264.7 cell and Human liver cancer tissue, SP20 cell using CD218b Polyclonal Antibody at dilution of 1:650 Calculated-MW:68 kDa	n, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Isotype IgG Purification Affinity purification Buffer Phosphate buffered solution Applications Recommended Dilut WB 1:500-1:2000 Data Image: Commende in the second in the se	
Purification Affinity purification Buffer Phosphate buffered solution Applications Recommended Dilut WB 1:500-1:2000 Data Image: Commended Dilut Western Blot analysis of RAW264.7 cell and Human liver cancer tissue, SP20 cell using CD218b Polyclonal Antibody at dilution of 1:650 Calculated-MW:68 kDa Preparation & Storage	
Buffer Phosphate buffered soluti Applications Recommended Dilut WB 1:500-1:2000 Data Western Blot analysis of RAW264.7 cell and Human liver cancer tissue, SP20 cell using CD218b Polyclonal Antibody at dilution of 1:650 Calculated-MW:68 kDa Preparation & Storage	
Applications Recommended Dilut WB 1:500-1:2000 Data Subscript of the second secon	
WB 1:500-1:2000 Data Image: Constraint of the second s	on
Data Data Data Data Data Data Data Data Data Data Solution Soluti	
Western Blot analysis of RAW264.7 cell and Human liver cancer tissue, SP20 cell using CD218b Polyclonal Antibody at dilution of 1:650 Calculated-MW:68 kDa Preparation & Storage	
Western Blot analysis of RAW264.7 cell and Human liver cancer tissue, SP20 cell using CD218b Polyclonal Antibody at dilution of 1:650 Calculated-MW:68 kDa Preparation & Storage	
$0.4 0.000 V_{-1} + 0.000$	
	nonths. Avoid freeze / thaw cycles.
temperature recommended	nonths. Avoid freeze / thaw cycles.
Background	-

Interleukin 18 receptor accessory protein, also known as IL18RAP and CDw218b (cluster of differentiationw218b), is a human gene. The protein encoded by this gene is an accessory subunit of the heterodimeric receptor for IL18. This protein enhances the IL18 binding activity of IL18R1 (IL1RRP), a ligand binding subunit of IL18 receptor. The coexpression of IL18R1 and this protein is required for the activation of NF-kB and MAPK8 (JNK) in response to IL18.

For Research Use Only