

# IL-8/CXCL8 Monoclonal Antibody(Capture)

catalog number: AN002140P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human IL-8/CXCL8 protein expressed by Mammalian
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG1
<b>Clone</b>	5G4
<b>Purification</b>	Protein A/G Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
----------------------	-----------

## Data



Sandwich ELISA-Recombinant Human IL-8/CXCL8 protein standard curve. Background subtracted standard curve using IL-8/CXCL8 antibody(AN002140P)(Capture), IL-8/CXCL8 antibody(AN002150P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human IL-8/CXCL8 protein is 7.81-500 pg/mL.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

Interleukin 8 (IL-8), also known as CXCL8, which is a member of the CXC chemokine family. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to inflammatory stimuli. IL-8 has two primary functions. It induces chemotaxis in target cells, primarily neutrophils but also other granulocytes, causing them to migrate toward the site of infection. IL-8 also induces phagocytosis once they have arrived. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. IL-8 is also known to be a potent promoter of angiogenesis. IL-8 has been associated with tumor angiogenesis, metastasis, and poor prognosis in breast cancer. IL-8 may present a novel therapeutic target for estrogen driven breast carcinogenesis and tumor progression.

## For Research Use Only