
Recombinant Human ERα protein (His Tag)

Catalog Number: PDEH101059

Note: Centrifuge before opening to ensure complete recovery of vial contents.

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Description	
Calculated MW32.9 kDaObserved MW38 kDaAccessionO75616Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Species	Human
Observed MW38 kDaAccessionO75616Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Human ERa protein Ala51-Asn350, with an C-terminal His
AccessionO75616Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	32.9 kDa
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	38 kDa
Properties Purity > 95% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Formulation Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Accession	O75616
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Bio-activity	Not validated for activity
Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Properties	
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 95% as determined by reducing SDS-PAGE.
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Shippingreconstituted samples are stable at < -20°C for 3 months.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Mannitol.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
	Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of		Mannitol.
it is reconstruction of a stock solution of	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.



> 95 % as determined by reducing SDS-PAGE.

Background

Data

Transports L-glutamate and also L-and D-aspartate. Essential for terminating the postsynaptic action of glutamate by rapidly removing released glutamate from the synaptic cleft. Acts as a symport by cotransporting sodium.