

Recombinant Human CD8 beta/CD8B Protein (Fc Tag)

Catalog Number:PKSH031296

 **DIA-AN®**
by Elabscience

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|--|
| Synonyms | T-Cell Surface Glycoprotein CD8 Beta Chain;CD8b;CD8B;CD8B1;LEU2;LYT3;LYT3;P37 |
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Met 1-Pro 170 |
| Accession | P10966-1 |
| Calculated Molecular Weight | 44.0 kDa |
| Observed molecular weight | 50 kDa |
| Tag | C-hFc |

Properties

| | |
|-----------------------|---|
| Purity | > 93 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 93 % as determined by reducing SDS-PAGE.

Background

CD8B (CD8b molecule); also known as P37 and LEU2; contains 1 Ig-like V-type (immunoglobulin-like) domain. The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen; acting as a coreceptor; and the T-cell receptor on the T lymphocyte recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains; or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. P37 gene encodes the CD8 beta chain isoforms. Multiple alternatively spliced transcript variants encoding distinct membrane

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD8 beta/CD8B Protein (Fc Tag)

Catalog Number:PKSH031296



associated or secreted isoforms have been described. A pseudogene; also located on chromosome 2; has been identified. CD8 is thought to play a role in the process of T-cell mediated killing.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com