

Recombinant DOPA Decarboxylase/DDC Monoclonal Antibody

catalog number: **AN301705L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Rat;Mouse
Immunogen	Recombinant human DOPA Decarboxylase / DDC fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A413
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution


WB	1:500-1:2000
IHC	1:200-1:1000
IF	1:50

Data




Western Blot with DOPA Decarboxylase/DDC Monoclonal Antibody at dilution of 1:2000. Lane 1: HepG2, Lane 2: Mouse adrenal gland

Observed-MW:53 kDa
Calculated-MW:53 kDa



Immunofluorescent analysis of (100% Ice-cold methanol) fixed PC-12 cells using anti-DOPA Decarboxylase/DDC Monoclonal Antibody at dilution of 1:50.



Immunohistochemistry of paraffin-embedded Human kidney using DOPA Decarboxylase/DDC Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

L-DOPA decarboxylase (DDC) is a pyridoxal 5-phosphate (PLP)-dependent enzyme that catalyzes the decarboxylation of L-DOPA to dopamine and the decarboxylation of L-5HTP to serotonin. DDC is also regarded as a general biomarker for neuroendocrine tumors.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0