Recombinant Human Contactin 2/CNTN2 Protein (His Tag)

Catalog Number: PKSH031598

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human Contactin 2/CNTN2 protein Met 1-Asn 1012, with an C-
	terminal His
Calculated MW	109 kDa
Observed MW	140 kDa
Accession	NP_005067.1
Bio-activity	Measured by the ability of the immobilized protein to support the adhesion of C6 Rat
	brain glial cells. When 5 x 10^4 cells/well are added to CNTN2-coated plates (0.8 µg/ml and 100 µl/well), approximately 40%-60% will adhere specifically after 60 minutes at 37°C.
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 $$
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.



> 95 % as determined by reducing SDS-PAGE.

Background

Data

Elabscience®

Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and NB-3. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. Contactin-2, also known as CNTN2, is a glycosylphosphatidylinositol (GPI)-anchored neuronal membrane protein that functions as a cell adhesion molecule. Contactin-2 is expressed by a subset of neuronal populations in the developing central nervous system (CNS) and peripheral nervous system (PNS). Contactin-2 is also expressed by oligodendrocytes and Schwann cells, which are myelinating glial cells of the CNS and PNS, respectively. Contactin-2 may play a role in the formation of axon connections in the developing nervous system. Contactin-2 is also involved in glial tumorigenesis and may provide a potential target for therapeutic intervention. During embryonic development, Contactin-2 interacts either in a homophilic, or heterophilic fashion with various transmembrane proteins.