## Recombinant Human SMAD4 Protein (His Tag)

## Catalog Number: PKSH033066

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human SMAD4 protein Met 1-Asp552, with an C-terminal His
Calculated MW	61.5 kDa
Observed MW	62 kDa
Accession	Q13485
Bio-activity	Not validated for activity
Properties	
Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.0.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

## Data



> 80 % as determined by reducing SDS-PAGE.

## Background

SMAD Family Member 4 (SMAD4) is a cytoplasmic protein that belongs to the Dwarfin/SMAD family. SMAD4 contains one MH1 (MAD homology 1) domain and one MH2 (MAD homology 2) domain. It is the component of the heterotrimeric SMAD2/SMAD3-SMAD4 complex that forms in the nucleus and is required for the TGF-mediated signaling. SMAD4 promotes binding of the SMAD2/SMAD4/FAST-1 complex to DNA and provides an activation function required for SMAD1 or SMAD2 to stimulate transcription. SMAD4 may act as a tumor suppressor. It positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator. Mutations or deletions in SMAD4 have been shown to result in pancreatic cancer, juvenile polyposis syndrome, and hereditary hemorrhagic telangiectasia syndrome.

For Research Use Only