

A Reliable Research Partner in Life Science and Medicine

## PE Anti-Mouse CD32 Antibody[S17012B]

Catalog Number: GFH00847D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Mouse Host Rat

IsotypeRat IgG2b, κClone No.S17012B

Isotype Control PE Rat IgG2b, κ Isotype Control[LTF-2] [Product GFH09842D]

Conjugation PE

**Conjugation Information** PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.


#### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### Data



Staining of C57BL/6 murine splenocytes cells with PE Anti-Mouse CD32 Antibody[S17012B] (left) or PE Rat IgG2b,  $\kappa$  Isotype Control (right). Total viable cells were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

#### **Antigen Information**

Alternate Names Fcr-2;Fcr-3;Ly-17;LyM-1;Lym-1;fcRll;FcgRll;fcgr2b;CD32抗体;CD32流式抗体;小鼠CD32;

Web: www.elabscience.cn

小鼠CD32抗体;小鼠CD32流式抗体;GFH00847

Uniprot ID P08101

#### For Research Use Only

# Elabscience®

### Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Gene ID Background 14130

CD32 (Fcgr2) is a 40 kD transmembrane glycoprotein, member of the immunoglobulin superfamily. The extracellular region of CD32 consists of two lg C-type domains that binds the Fc region from monomeric lgG with low affinity, but binds immune complexes efficiently. CD32 can mediate phagocytosis of immune complexes and modulate cell activation. CD32 is expressed by Macrophages, neutrophils, mast cells and B cells.