

Recombinant Mouse sFRP4 Protein (His Tag)

Catalog Number: PKSM040882

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse sFRP4 protein Met 1-Ser 351, with an C-terminal His
Calculated MW	39.4 kDa
Observed MW	55-60 kDa
Accession	NP_057896.1
Bio-activity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.6

SFRP family consists of five secreted glycoproteins in humans acting as extracellular signaling ligands. Each is approximately 300 amino acids in length and contains a cysteine-rich domain (CRD) that shares 30-50% sequence homology with the CRD of Frizzled (Fz) receptors, a putative signal sequence, and a conserved hydrophilic carboxy-terminal domain. SFRPs act as soluble modulators of Wnt signaling, counteracting Wnt-induced effects at high concentrations and promoting them at lower concentrations. SFRPs are able to bind Wnt proteins and Fz receptors in the extracellular compartment. The interaction between SFRPs and Wnt proteins prevents the latter from binding the Fz receptors. The Wnt pathway plays a key role in embryonic development, cell differentiation and cell proliferation. SFRP4 is a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins called FZ domain and a NTR domain. Mouse SFRP4 is highly expressed in the ovary, and is localized to granulosa cells of periovulatory follicles and corpora lutea. It plays a critical role in placental development and implantation, and is also an important factor in the development of the decidual fibrinoid zone, and in trophoblast apoptosis.