

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Red 780 Anti-Human CD86 Antibody[BU63]

Catalog Number: E-AB-F1012S

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. BU63

Isotype Control Elab Fluor® Red 780 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792S]

**Conjugation** Elab Fluor<sup>®</sup> Red 780

**Conjugation Information** Elab Fluor<sup>®</sup> Red 780 is designed to be excited by the Red (627-640 nm) laser and

detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter).


**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

## Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Staining of normal human peripheral blood cells with Anti-

Human CD14 FITC and Anti-Human CD86 Elab Fluor<sup>®</sup> Red 780 (left) or Mouse IgG1, κ Isotype Control Elab Fluor<sup>®</sup> Red 780 (right). Cells in the monocyte gate were used for analysis.

### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

### **Antigen Information**

Alternate Names Activation B7-2 antigen;Cd86;ETC-1;Early T-cell costimulatory molecule 1;T-lymphocyte

activation antigen CD86

Uniprot ID P42081

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 942

CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, monocytes/macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is the ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce costimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can bind to CD152 as well, also known as CTLA-4, to deliver an inhibitory signal to T cells.