Elabscience®


Elab Fluor[®] Violet 450 Anti-Mouse TCRβ Antibody[H57-597]

Catalog Number: E-AB-F1123Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Armenian Hamster
lsotype	Armenian Hamster IgG
Clone No.	H57-597
Isotype Control	Elab Fluor [®] Violet 450 Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with Elab Fluor[®] Violet 450 Anti-Mouse TCR β Antibody (filled gray histogram)

or Elab Fluor[®] Violet 450 Armenian hamster IgG Isotype Control (empty black histogram).

Preparation & Storage	•
Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names Gene ID	TCR-ββ-TCR;TCR-β chain 21577

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>www.elabscience.com</u>

Elabscience®

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

Background

T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ and a δ chain (TCR γ/δ). TCR- β is a member of the immunoglobulin superfamily and a component of the CD3/TCR complex (along with TCR- α). It is expressed on α/β TCR-bearing T cells and thymocytes. The CD3/TCR complex plays a key role in antigen recognition, signal transduction, and T cell activation.