

## Recombinant Human G-CSFR/CD114 Protein (His Tag)

**Catalog Number:** PKSH031748

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human G-CSFR/CD114 protein Met 1-Pro 621, with an C-terminal His
<b>Calculated MW</b>	68.0 kDa
<b>Observed MW</b>	92 kDa
<b>Accession</b>	NP_000751.1
<b>Bio-activity</b>	Measured by its ability to inhibit the GCSF-induced proliferation of NFS60 mouse myeloid cells. The ED <sub>50</sub> for this effect is typically 50-250 ng/mL in the presence of 0.125ng/mL of recombinant human GCSF.

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Granulocyte Colony Stimulating Factor Receptor (G-CSFR), also known as CD114, which belongs to the cytokine receptor superfamily, is a cell surface receptor for colony stimulating factor 3 (CSF3). It is a critical regulator of granulopoiesis. This type I membrane protein has a composite structure consisting of an immunoglobulin(Ig)-like domain, a cytokine receptor-homologous (CRH) domain and three fibronectin type III (FNIII) domains in the extracellular region. Mutations in the G-CSF receptor leading to carboxy-terminal truncation transduce hyperproliferative growth responses, and are implicated in the pathological progression of severe congenital neutropenia (SCN) to acute myelogenous leukemia (AML). Additionally, autocrine/paracrine stimulation of G-CSFR may be important in the biology of solid tumors, including metastasis.