Recombinant Human GDF-15 protein (His Tag)

Catalog Number: PDEH101053

Note: Centrifuge before opening to ensure complete recovery of vial contents.

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Description			
Calculated MW17.3 kDaObserved MW20 kDaAccessionQ99988Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Species	Human		
Observed MW20 kDaAccessionQ99988Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Human GDF-15 protein Gln151-Ile308, with an C-terminal His		
AccessionQ99988Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	17.3 kDa		
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	20 kDa		
PropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Accession	Q99988		
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Bio-activity	Not validated for activity		
Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Properties			
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	>95% as determined by reducing SDS-PAGE.		
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method		
Shippingreconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Mannitol.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
	Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%		
		Mannitol.		
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
0.5 mg/mL. Concentration is measured by UV-Vis.		0.5 mg/mL. Concentration is measured by UV-Vis.		

Data

KDa	М	R
135 100	-	
75	-	
65	1	
45	-	
35	-	
25		•
15		

> 95 % as determined by reducing SDS-PAGE.

Background

Growth/differentiation factors (GDF-1 to GDF-15) are members of the BMP family of TGF-beta superfamily proteins. They are produced as inactive preproproteins which are then cleaved and assembled into active secreted homodimers. GDF dimers are disulfide-linked with the exception of GDF-3 and-9. GDF proteins are important during embryonic developmen t, particularly in the skeletal, nervous, and muscular systems.