

AF/LE Purified Anti-Human CD147 Antibody[HIM6]

catalog number: E-AB-F10560

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD147 protein |
| Host | Mouse |
| Isotype | Mouse IgG1, κ |
| Clone | HIM6 |
| Purification | >98%, Protein A/G purified |
| Conjugation | None (AF/LE) |
| Buffer | Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method |

Applications

Recommended Dilution

| | |
|------------|---|
| FCM | 2 $\mu\text{g/mL}$ (1×10^5 - 5×10^5 cells) |
|------------|---|

Data



Human peripheral blood lymphocytes were stained with 0.2 μg AF/LE Purified Anti-Human CD147 Antibody[HIM6] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions. |
| Shipping | Ice bag |

Background

CD147, also known as neurothelin or basigin, is a member of the Ig superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.

For Research Use Only