## Recombinant Human KIR2DL1/CD158a Protein (Fc Tag)

## Catalog Number: PKSH030712

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human KIR2DL1/CD158a protein Met 1-His 245, with an C-
	terminal hFc
Calculated MW	51.7 kDa
Observed MW	70 kDa
Accession	NP_055033.2
<b>Bio-activity</b>	Not validated for activity
Properties	
Purity	>90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	
	KDa MK R



> 90 % as determined by reducing SDS-PAGE.

Background

## **Elabscience**®

Killer cell immunoglobulin-like receptor 2DL1 or KIR2DL1 is an inhibitory natural Killer cell immunoglobulin-like receptor with two extracellular immunoglobulin domains. KIR2DL1 is a member of the Killer cell immunoglobulin-like receptor family whose members are classified by the number of the extracellular immunoglobulin domains and the length of the cytoplasm domain. KIR2DL1 is a transmembrane glycoprotein expressed by natural killer cells and subsets of T cells. KIR2DL1 down-regulates the cytotoxicity of NK cells upon recognition of specific class I major histocompatibility complex (MHC) molecules on target cells. It has been reported that the KIR2DL1 bound to its class I MHC ligand; HLA-Cw4. The KIR2DL1-HLA-Cw4 interface exhibits charge and shape complementarity. Specificity is mediated by a pocket in KIR2DL1 that hosts the Lys80 residue of HLA-Cw4. Many residues conserved in HLA-C and in KIR2DL receptors make different interactions in KIR2DL1-HLA-Cw4 and in a previously reported KIR2DL2-HLA-Cw3 complex. A dimeric aggregate of KIR-HLA-C complexes was observed in one KIR2DL1-HLA-Cw4 crystal.