A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse CD79A/Ig-alpha Protein (ECD, Fc Tag)

Catalog Number: PKSM040512

Note: Centrifuge before opening to ensure complete recovery of vial contents.

# Description

Species Mouse

Source HEK293 Cells-derived Mouse CD79A/Ig-alpha protein Met1-Arg137, with an C-

terminal hFc

 Calculated MW
 39.0 kDa

 Accession
 NP\_031681.2

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per  $\mu$ g of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



> 90 % as determined by reducing SDS-PAGE.

# Background

Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B-cells.

#### For Research Use Only