# **BLNK Polyclonal Antibody**

catalog number: E-AB-18540





Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human BLNK
Host	Rabbit
Is otype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:40-1:200

#### Data






Western blot analysis of LOVO cell lysate using BLNK

### Polyclonal Antibody at dilution of 1:600

#### **Observed-MV:Refer to figures**

#### Calculated-MV:50 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using BLNK Polyclonal Antibody at dilution of 1:40(×200)

Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using BLNK Polyclonal Antibody at dilution of  $1:40(\times 200)$ 

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

#### Background

## For Research Use Only

# **BLNK Polyclonal Antibody**

#### catalog number: E-AB-18540



This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia.

### For Research Use Only