

A Reliable Research Partner in Life Science and Medicine

# FITC Anti-Mouse CD5 Antibody[53-7.3]

Catalog Number: E-AB-F1185C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ Clone No. 53-7.3

FITC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832C] Isotype Control

Conjugation

**Conjugation Information** FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

## **Applications**


#### Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### **Data**



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD3 Antibody and FITC Anti-Mouse CD5 Antibody (Left). Splenocytes are stained with APC Anti-Mouse CD3 Antibody and FITC Rat IgG2a, κ Isotype Control (Right).

## **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

# **Antigen Information**

**Alternate Names** CD5;Cd5;Ly-1;Lymphocyte antigen 1;Lyt-1

**Uniprot ID** P13379 Gene ID 12507

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web:www.elabscience.com Email:techsupport@elabscience.com

Rev. V1.4



#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Background

CD5 is a 67 kD protein, also known as Lyt-1, Ly-1, T1, Tp67, or Ly-12. It is a member of the scavenger receptor cysteine-rich protein superfamily (SRCR) and primarily expressed on thymocytes, T cells, and B-1 cells. Although mature  $\alpha/\beta$  T cells express high levels of CD5, very few  $\gamma/\delta$  T cells express this antigen. The interaction of CD5 with CD72, gp35-37, TCR, or BCR is involved in T and B cell activation.

Fax: 1-832-243-6017