Elabscience®


Elab Fluor[®] Violet 450 Anti-Mouse CD4 Antibody[RM4-5]

Catalog Number: E-AB-F1353UQ

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat lgG2a, κ
Clone No.	RM4-5
Isotype Control	Elab Fluor [®] Violet 450 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833Q]
Conjugation	Elab Fluor [®] Violet 450
Conjugation Information	Elab Fluor [®] Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended

concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].



C57BL/6 murine splenocytes are stained with FITC Anti-

Mouse CD8a Antibody and Elab Fluor[®] Violet 450 Anti-Mouse CD4 Antibody[RM4-5] (Left). Splenocytes are stained

with FITC Anti-Mouse CD8a Antibody and Elab Fluor[®] Violet 450 Rat IgG2a, κ Isotype Control (Right).

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	L3T4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4;T4
Uniprot ID	P06332

For Research Use Only Toll-free: 1-888-852-8623

Web:www.elabscience.com

Data

Elabscience®

Gene ID Background

12504

CD4 is a 55 kD protein also known as L3T4 or T4. It is a member of the Ig superfamily, primarily expressed on most thymocytes and a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a co-receptor with the TCR during T cell activation and thymic differentiation by binding MHC class II and associating with the protein tyrosine kinase lck.